

BR ACTIVATOR®



Product Introduction:

Active Ingredient	Content & Formulation
Brassinolide	0.01%SL, 0.01%SP

Product Introduction:


It is used as soluble concentrate for a variety of crops and a lot of orchard and fruit products. It can result in remarkable promotion of crops growth and productivity; and a promotion of ability of crops disease-resistant, salt-resistant and cold-resistant to alleviate the harmfulness caused by using of herbicides.



Advantage:

1. Promote growth, increase yield, and improve quality.
2. Promote the percentage of fruit setting and fruit's enlarging.
3. Increase seed setting percentage rate and fruits thousand kernel weight.
4. Improve the ability of cold resistance, drought resistance, and salt soil resistance, enhance growth recovery ability after flooding.
5. Improve the ability of disease resistance.
6. Regulate the differentiation among the plant.
7. Efficiently relieve harmful effect of pesticides, fungicides, and herbicides.
8. Improve the activity and vitality of seeds and germination percentage.
9. Efficiently prolong the duration time of natural preservation and transportation of fruits, vegetables and flowers when applied 5-10 days before harvest.

Crops: Rice, Cotton, Corn, Tea, Water melon, Banana, Wax apple, Grape, Peach, Pear, Cucumber, Chili.



Rice



Peach



Pear



Grapes



Cucumbers

Uses and Recommendations:

Crops	Effectiveness	Dosage	Application method
Rice/Cereals	Increase germination	5gram-6gram with 15L water	Seeds soaking treatment or seeds treatment
	Promote crop growth	4.5gram-5gram with 15L water	Spray
Corn/Maize	Increase germination	5gram-5.5gram with 15L water	Seeds soaking treatment or seeds treatment
	Promote crop growth	5gram-6gram with 15L water	Spray
Melons/Vegetables	Increase germination	4gram-5gram with 15L water	Seeds soaking treatment or seeds treatment
	Promote crop growth	5gram-5.5gram with 15L water	Spray
Cotton	Promote crop growth	4.5gram-5.0gram with 15L water	Spray
Tea/Chili/Fruits	Promote crop growth	4gram-5gram with 15L water	Spray

Cautions:

1. Do not spray on rainy days, re-spray if it rains within 6h after spraying.
2. Pay attention to wind direction changes, don't spray against the wind, but follow along the winds direction to avoid inhalation.
3. Do not store near seeds food and other feedstuffs.
4. Do not re-use container, do not casually discarded. Dispose product containers, waste containers, residues according local health and environmental regulations.
5. Wear protective clothing and gloves, avoid inhaling liquid when using this product, do not eat or drink during application, wash hands and wash face timely after application.